

**An automated analysis tool for the classification
of Sea Surface Temperature imagery**

Marco Reggiannini^a, Oscar Papini^a, Gabriele Pieri^a

^aInstitute of Information Science and Technologies,

National Research Council of Italy,

Via G. Moruzzi 1, 56124 Pisa, Italy

email: {name.surname}@isti.cnr.it

Abstract— Sea observation through remote sensing technologies plays an essential role in understanding the health status of the marine coastal environment, its fauna species and their future behaviour. Accurate knowledge of the marine habitat and the factors affecting faunal variations allows us to perform predictions and adopt proper decisions. This paper concerns the proposal of a classification system devoted to recognising marine mesoscale events. These phenomena are studied and monitored by analysing Sea Surface Temperature imagery. Currently, the standard way to perform such analysis relies on experts manually visualising, analysing, and tagging large imagery datasets. Nowadays, the availability of remote sensing data has increased so much that it is desirable to replace the labour-intensive, time-consuming and subjective manual interpretation with automated analysis tools. The results presented in this work have been obtained by applying the proposed approach to images captured over the southwestern region of the Iberian Peninsula.

Keywords: Image Processing; Remote Sensing; Mesoscale Patterns; Sea Surface Temperature; Machine Learning; Climate Change

1 INTRODUCTION

To achieve a broader understanding and evaluation of the sea environment, an improvement in marine observation is required. Among all the relevant underlying processes in such a differentiated biological system, mesoscale events such as upwelling, countercurrents and filaments are of particular interest and constitute the subject of our analysis. These events, which transport deeper, colder and nutrient-rich waters to the surface, and affect the biological parameters of the habitat, enhancing the local biodiversity [7], can be observed by analysing Sea Surface Temperature (SST) recorded in remote sensing imagery.

Identifying and categorising upwelling regimes occurring in a marine ecosystem is an essential achievement for its characterisation. The main objective of this paper is to propose a method for performing an automatic classification of images in place of the usual manual one completed by experts. When the number of images approaches the thousands, i.e. the typical order of magnitude having the goal to investigate long term and climate-related changes, the manual procedure is not manageable anymore. The method is applied to the Iberia/Canary Current System (ICCS), one of the least studied among the upwelling ecosystems [1]. Despite a general circulation similar to others, in ICCS we have diverse factors having a profound impact on the whole region.

The method proposed in this work is based on implementing an automatic procedure for classifying large datasets of images according to the different regimes of observable upwelling patterns. Such classification consists of several

stages: starting from the extraction of quantitative features from a region of interest in the SST maps, proceeding to the characterisation of specific temperature patterns, which are correlated with the water flows between geographical points at different temperatures. The latter stage is performed by applying a set of rules to the computed features, which enable the assignment of a final class label to the considered region. This method follows and completes the preliminary analysis performed in [4] and [6].

The paper is arranged as follows: Section 2 provides a description of the employed dataset and the related ground truth classification; Section 3 reports on the pipeline used in our methods and describes a study case; Section 4 concludes the paper by discussing the outcomes of this work and providing a few considerations about future perspectives.

2 MATERIALS

For the purposes of this work, SST data captured by Metop-A/B (EUMETSAT) and Aqua (NASA) have been collected and processed. Only data covering the region of interest were downloaded for each source (whose respective details are reported in Table 1). In particular, points with latitude between 35° and 40° N and longitude between 12° and 6° W were considered, resulting in 2–3 images per day at most.

Table 1. Data specifications

Satellite	Sensor Type	Spatial Resolution (km)	Temperature Resolution ($^\circ\text{C}$)
Metop-A/B [3]	AVHRR	1	10^{-2}
Aqua [2]	MODIS	1	$5 \cdot 10^{-3}$

Expert oceanographers have preliminarily inspected the collected data to identify recurring SST patterns based on the detection of relevant mesoscale features (water filaments, upwelling jets and countercurrents). This way, it was possible to identify four prevailing patterns, named E1–E4 (see [6] for a detailed description). Furthermore, each image was labelled according to the observed pattern, returning a ground truth dataset that could be used as a reference for the classifier implementation.

3 SST ANALYSIS

In order to better analyse the different types of upwelling patterns, SST data are retrieved from the sources described in the previous section and arranged in a *spaghetti plot*, which is a simultaneous representation of the different SST trends for a given geographical area and a time interval. It is obtained by first dividing the considered area into a grid of small squares (whose size may be equal to or larger than the image spatial resolution). Then, for each square, the SST spatial average value is computed for each time sample in the dataset falling within the considered time window. Finally, the obtained ensemble of averaged SSTs is plotted versus time within the same diagram.

Figure 1 shows an example of an event classified as E4 in the ground truth and the spaghetti plots corresponding to the selected areas. Events of type E4 are characterised by the presence of a warm countercurrent originating in the Gulf of Cádiz and running along the southern Iberian coast, eventually reaching Cape St. Vincent (see Figure 1c). A cold water filament going westwards is also recognisable (see Figure 1b), which is a pattern typical for events of type E1. In this case, the squares' size and the time interval are 0.25° and 15 days respectively (notice that the ground truth event occurs at the end of the time

window). After several tests, these specific values have been chosen since they return a better agreement between the results and the ground truth.




Figure 1. Event of 7 October 2017 at around 21:00 UTC. (a) SST map at the date of the event; (b) detail of the SST in the reference area for spaghetti plot I (latitude between 37.25° and 38° N, longitude between 10.75° and 10° W); (c) detail of the SST in the reference area for spaghetti plot II (latitude between 36.5° and 37.25° N, longitude between 8.75° and 8° W); (d) reference grid for both plots (dimension of squares 0.25°); (e,f) generated spaghetti plots.

A spaghetti plot is then processed to extract statistical features, which depend on the SST signal in each square and its neighbourhood. These features

are later used to classify the considered area, which is then associated with one of the four mesoscale patterns.

Let a be a square in the grid. As said, we have a temporal series of spatial SST averages in a , say μ_i , computed at times t_i , $i = 1, \dots, n$. Notice that n may change from square to square, since it depends on the number of SST values captured by the sensor. In fact, the SST recording may fail for some parts of the area of interest (e.g. due to interfering clouds disturbances). Because of these considerations, the number of samples n can be considered as an index of reliability for the classification of the square a . The statistics features computed for a are:

1. the temporal mean $\mu(a)$, defined as the mean of the values μ_i ;
2. the standard deviation $\sigma(a)$, defined as the standard deviation of the values μ_i ;
3. the linear regression coefficient $\theta(a)$, defined as the slope of the straight line that better interpolates the values (t_i, μ_i) .

The values μ , σ and θ are computed for every square in the grid. Then, a set of rules is applied to obtain, for each square a , an array of four scores (e_1, e_2, e_3, e_4) , with $e_j \in [0,1]$. The value e_j represents a belief index for the event of type E_j to have occurred inside a at the end of the considered time interval. The implementation of the rules is a crucial component for the classifier. Indeed, they are handcrafted so that the score e_j is boosted only if the behaviour of the features μ , σ and θ , inside and in the neighbourhood of the square a , matches the one observed in the case of an E_j pattern.

The classification of a square is finally completed by considering the maximum score $e_m = \max\{e_1, e_2, e_3, e_4\}$: if e_m is above a certain threshold,

empirically defined, then the square is labelled “Em”; otherwise no label is assigned. Figure 2 represents a heatmap with the classification results to the event of Figure 1, with each square coloured with the corresponding classification label. Also, each square is labelled with the numerical percentage of the related SST data, which is proportional to the n value, as discussed above.




Figure 2. Labels given to each square of the grid, depending on their scores.

4 DISCUSSION AND CONCLUSION

In this work, a methodology for classifying of upwelling events based on the analysis of SST time series has been proposed. Preliminary tests proved that the proposed method succeeds in classifying different mesoscale events. A few considerations can be pointed out concerning the presented case study (Figure 1). First, it is worth remarking that the labelling returned by the classifier agrees with the ground truth: among the squares located in the area where E4 events usually occur, those that fulfilled the previously mentioned data abundance constraints have been labelled accordingly (Figure 2). Second, the proposed method extracts features that, as previously discussed, take into account not only the SST final observation, corresponding to the ground truth

label, but also the SST variations captured in the preceding time window. This is the reason behind the presence of squares classified differently from E4, in apparent conflict with the ground truth. Since the proposed approach considers the SST signal over an extended range of time, it is reasonable that more than one label is assigned, in agreement with the multiple observed mesoscale events. It is even more so considering that inside the presented case study's dataset, different ground truth labels have been assigned to images captured very close in time. For example, on 6th October, two distinct events are observed: one classified as E1 in the ground truth approximately at 10:00 UTC, and a second one around 21:20 UTC classified as E4.

The test and validation of the proposed algorithm are carried out and will continue as part of the activities of the EU H2020 project NAUTILOS [5].

ACKNOWLEDGEMENTS

The authors express gratitude to Prof. Flávio Martins and Dr João Janeiro from the University of Algarve, Centre for Marine and Environmental Research, for their support.

FUNDING

This paper is part of a project that has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 101000825 (NAUTILOS).

CONFLICT OF INTERESTS

The process of writing and the content of the article does not give grounds for raising the issue of a conflict of interest.

COMPLIANCE WITH ETHICAL STANDARDS

This article is a completely original work of its authors; it has not been published before and will not be sent to other publications until the PRIA editorial board decides not to accept it for publication.

REFERENCES

1. F. P. Chavez and M. Messié, “A comparison of Eastern Boundary Upwelling Ecosystems,” *Progress in Oceanography* 83 (1), 80–96 (2009). <https://doi.org/10.1016/j.pocean.2009.07.032>
2. NASA/JPL, *GHRSSST Level 2P Global Sea Surface Skin Temperature from the Moderate Resolution Imaging Spectroradiometer (MODIS) on the NASA Aqua satellite (GDS2)* (2020). <https://doi.org/10.5067/GHMDA-2PJ19>
3. OSI SAF, *Full resolution L2P AVHRR Sea Surface Temperature MetaGRanules (GHRSSST) – Metop* (2011). https://doi.org/10.15770/EUM_SAF_OSI_NRT_2013
4. O. Papini, M. Reggiannini, and G. Pieri, “SST Image Processing for Mesoscale Pattern Identification,” *Eng. Proc.* 8, 5 (2021). <https://doi.org/10.3390/engproc2021008005>
5. G. Pieri et al., “New technology improves our understanding of changes in the marine environment,” *Proc. of the 9th EuroGOOS International Conference* (2021).
6. M. Reggiannini, J. Janeiro, F. Martins, O. Papini, and G. Pieri, “Mesoscale Patterns Identification Through SST Image Processing,” *Proc. of the 2nd*

International Conference on Robotics, Computer Vision and Intelligent Systems, 165–172. <https://doi.org/10.5220/0010714600003061>

7. R. Varela, F. P. Lima, R. Seabra, C. Meneghesso, and M. Gómez-Gesteira, “Coastal warming and wind-driven upwelling: A global analysis,” *Science of The Total Environment* 639, 1501–1511 (2018). <https://doi.org/10.1016/j.scitotenv.2018.05.273>

AUTHORS PROFILE



Marco Reggiannini (Viareggio, Italy, 1981), M.Sc. (2008) in Physics and Ph.D. (2016) in Automation Engineering from the University of Pisa, currently technologist at the Signals and Images Laboratory at ISTI-CNR, Pisa. His main interests concern signals and images processing, scene understanding, pattern recognition and machine learning.



Oscar Papini (Lecco, Italy, 1989), Ph.D. (2018) in Mathematics from the University of Pisa, currently post-doctoral fellow at the Signal and Images Laboratory at ISTI-CNR, Pisa. His main interests are machine learning and mathematical methods for image analysis.



Gabriele Pieri (Pescia, Italy, 1974), M.Sc. (2000) in Computer Science from the University of Pisa, since 2001 joined the Signals and Images Laboratory at ISTI-CNR, Pisa, working in the field of image analysis. His main interests include neural networks, machine learning, industrial diagnostics and environmental monitoring. He is author of more than twenty papers.